

**PROOF OF A SPECIAL CASE OF THE YELTON-GAINES CONJECTURE ON ISOMORPHIC
DESSINS**

Claudia Raithel¹, R. Perlis^{*2}, Univeristy of Michigan- Ann Arbor¹, Department of Mathematics,
Ann Arbor, MI 48109, Louisiana State University², Department of Mathematics, Baton Rouge, LA
70803, perlis@math.lsu.edu

Let (ρ_0, ρ_1) and (ρ'_0, ρ'_1) be two ordered pairs of permutations in S_n and let t be a divisor of n . The Yelton-Gaines conjecture states that if at least one of these four permutations is a product of n/t disjoint t -cycles, and if there is a *strong* isomorphism (definition below) $\phi : \langle \rho_0, \rho_1 \rangle \rightarrow \langle \rho'_0, \rho'_1 \rangle$ between the two subgroups of S_n generated by the elements in each ordered pair, then there is a fixed permutation τ in S_n that simultaneously conjugates ρ_i to ρ'_i for $i = 0, 1$. The conclusion of this conjecture can be restated to say that the two *dessins d'enfants* corresponding to the two ordered pairs are isomorphic.

A proof of this conjecture is given in the case in which all of the initial four permutations are fixed-point-free involutions.